



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\log x_j \begin{vmatrix} 2 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 2 & 1 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 2 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 2 \end{vmatrix} = \begin{vmatrix} 2 & 1 & 0 & \cdots & \log a_1 & 0 & \cdots & 0 \\ 0 & 2 & 1 & \cdots & \log a_2 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \log a_j & 1 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & 0 & 0 & \cdots & \log a_n & 0 & \cdots & 2 \end{vmatrix}.$$

Denoting the determinant in the left member of the last equation by D and the one in the right member by A_j , we have

$$\log x_j = \frac{A_j}{D},$$

or

$$x_j = e^{\frac{A_j}{D}} \quad (j = 1, 2, \cdots, n).$$

Also solved by the PROPOSER, who used logarithms as in the last solution, though he did not use determinants.


GEOMETRY.

424. Proposed by H. E. TREFETHEN, Colby College.

In a given triangle ABC , determine by geometric demonstration the point O such that the sum of the distances, $AO + BO + CO$, shall be a minimum.

SOLUTION BY M. E. GRABER, Heidelberg University.

On each of the three sides of the triangle ABC , describe a segment of a circle to contain an angle of 120° . The arcs of these circles intersect at a point O , about which the angles are each 120° . Then $AO + BO + CO$ is a minimum. Suppose $OA > OB$ or OC . Using OA as a radius describe a circumference. Produce OC and OB to meet the circumference at M and N . Let L be any other point in ABC .



Draw LC , LM , LB , LN , AM , MN , NA . Then $AM = MN = NA$ and $LA + LN + LM > OA + ON + OM$. In order to prove this relation, it is evidently sufficient to prove that $OA + ON + OM <$ the sum of the distances from S (the vertex of an isosceles Δ on MN with the same altitude as ΔLMN)

to A , M , and N . With MS and NS as radii describe arcs cutting MO and NO in T and V . Join T and S , V and S , by straight lines. Then $\angle OVS$ is obtuse and therefore $\angle OSV$ is $< 30^\circ$. When $\angle SOV = 60^\circ$ and $\angle OSV = 30^\circ$, $OV = \frac{1}{2}OS$. When $\angle OSV < 30^\circ$, $OV < \frac{1}{2}OS$ and $OV + OT < OS$. Hence $SM + SN + SA > AO + OM + ON$ and $OM + ON + OA < LM + NL + LA$. Since $LC + CM > LM$ and $LB + BN > LN$, we have $LC + CM + LB + BN + LA > OA + OB + BN + OC + CM$. Hence, $LA + LB + LC > OA + OB + OC$.

Also solved by C. N. SCHMALL, DAVID L. MACKEY, A. M. HARDING and HARVY ROESER.

CALCULUS.

336. Proposed by EVA S. MAGLOTT, Ada, Ohio.

If a right cone stands on an ellipse, prove that its superficial area is $\frac{\pi}{2}(OA + OA')(OA \cdot OA')^{\frac{1}{2}}$ times $\sin \alpha$, where O is the vertex of the cone, A and A' extremities of the major axis of the ellipse, and α is the semi-angle of the cone.


SOLUTION BY ELMER SCHUYLER, Brooklyn, N. Y.

The formula for the surface $AA'LT$ is

$$\frac{\pi}{R-r} \sqrt{a^2 + (R-r)^2} \{R^2 - \frac{1}{2}(R+r) \sqrt{Rr}\}.$$

(See Finkel's *Mathematical Solution Book*, pp. 318-319).

Here $r = OA \sin \alpha$, $R = OA' \sin \alpha$, and $a = (OA' - OA) \cos \alpha$.



For brevity, put $OA = K$, $OA' = K'$. Then this formula becomes, after slight reduction,

$$\frac{\pi}{\sin \alpha} \cdot \sin^2 \alpha \{K'^2 - \frac{1}{2}(K + K') \sqrt{KK'}\} = \frac{\pi}{2} \sin \alpha \{2K'^2 - (K + K') \sqrt{KK'}\}.$$